



ELSEVIER

REVIEW ARTICLE

JOURNAL of  
CARDIOLOGY

Official Journal of the Japanese College of Cardiology

www.elsevier.com/locate/jjcc

# Beneficial effects of ezetimibe-based therapy in patients with dyslipidemia

Shin-ichiro Miura (Ph.D., M.D.)\*, Keijiro Saku (Ph.D., M.D.)

Department of Cardiology, Fukuoka University School of Medicine, Fukuoka, Japan

Received 6 March 2008; received in revised form 7 May 2008; accepted 9 May 2008  
Available online 30 June 2008

## KEYWORDS

Dyslipidemia;  
Low-density lipoprotein  
cholesterol;  
Ezetimibe;  
Niemann-Pick C1-like 1

**Summary** Treatment of dyslipidemia is important for the primary and secondary prevention of cardiovascular events. Although statins induce the intensive lowering of low-density lipoprotein (LDL) cholesterol levels, two-thirds of cardiovascular events are not prevented. Ezetimibe has been shown to be a selective inhibitor of the Niemann-Pick C1-like 1 (NPC1L1) transporter of cholesterol across the intestinal wall. Ezetimibe-based therapy may hold the promise of more intensive lowering of LDL cholesterol. This review will address the beneficial effects of ezetimibe in patients with dyslipidemia.

© 2008 Japanese College of Cardiology. Published by Elsevier Ireland Ltd. All rights reserved.

## Contents


Introduction.....	1
Ezetimibe monotherapy.....	2
Ezetimibe/statin combination therapy.....	3
Ezetimibe or ezetimibe/statin in combination with fibrate therapy.....	4
Genetic studies on ezetimibe therapy.....	5
Safety of ezetimibe.....	5
Conclusions.....	5
References.....	5

## Introduction

Ezetimibe (1-(4-fluorophenyl)-(3R)-[3-(4-fluorophenyl)-(3S)-hydroxypropyl]-(4S)-(4-hydroxypropyl)-2-azetidinone) is a potent inhibitor of cholesterol and phytosterol uptake [1]. Since low cholesterol absorption has been associated with


\* Corresponding author. Department of Cardiology, Fukuoka University School of Medicine, 7-45-1 Nanakuma, Jonan-Ku, Fukuoka 814-0180, Japan. Tel.: +81 92 801 1011; fax: +81 92 865 2692.

E-mail address: miuras@cis.fukuoka-u.ac.jp (S.-i. Miura).



**Figure 1** Unadjusted Kaplan–Meier curve for total mortality in the lowest quartile of serum cholestanol-to-cholesterol ratio ( $n=94$ ) vs. other quartiles combined ( $n=284$ ).

a lower rate of total mortality [2] (Fig. 1), pharmacological intervention through the inhibition of intestinal cholesterol absorption may be a useful strategy for treating patients with dyslipidemia and/or cardiovascular disease. Niemann-Pick C1-like 1 (NPC1L1) has recently been identified, and has been shown to have features of a plasma membrane transporter, including a secretion signal, 13 predicted transmembrane domains, extensive N-linked glycosylation sites and a sterol-sensing domain [3] (Fig. 2). It is highly expressed on the surface of absorptive jejunal enterocytes. NPC1L1 has been shown to be a direct target of ezetimibe, and an ezetimibe-sensitive pathway plays a role in intestinal cholesterol absorption [4]. Despite the availability of statins, many patients do not achieve lipid targets. Combination therapy with statins and ezetimibe that act via a complementary pathway may allow additional patients to achieve their recommended cholesterol goals. These findings represent an exciting new area in the treatment of dyslipidemia. Ezetimibe-based therapy may hold the promise of more intensive lowering of low-




**Figure 3** Changes in serum lipid profile in placebo ( $n=70$ , black bars), ezetimibe therapy ( $n=61$ , white bars), simvastatin therapy ( $n=263$ , gray bars), and ezetimibe+simvastatin therapy ( $n=274$ , dotted bars) in patients with primary hypercholesterolemia. LDL, low-density lipoprotein; HDL, high-density lipoprotein; TG, triglyceride. \* $p < 0.05$  vs. simvastatin therapy.

density lipoprotein (LDL) cholesterol. This review will address the beneficial effects of ezetimibe in patients with dyslipidemia.

### Ezetimibe monotherapy

Ezetimibe can effectively lower plasma LDL cholesterol to decrease cholesterol absorption [5]. Ezetimibe at 10mg/day induced an about 20% reduction in LDL cholesterol [6] (Fig. 3). In addition, ezetimibe decreased triglyceride (TG) by about 8% and increased high-density lipoprotein (HDL) cholesterol by about 5%. In patients with primary dyslipidemias, ezetimibe (10mg/day)



**Figure 2** Secondary structure of NPC1L1.

therapy for 16 weeks reduced total, LDL and non-HDL cholesterol values as well as apolipoprotein B concentrations [7]. Patients with TG values >150 mg/dL had significantly greater reductions in the concentrations of small, dense LDL particles compared to those with normal TG levels (49% vs. 19%, respectively;  $p < 0.05$ ). With respect to individual LDL subfractions, cholesterol was significantly reduced by ezetimibe in cholesterol in nearly all LDL subfractions [8]. These results indicate that ezetimibe significantly reduces LDL cholesterol by inhibiting the reabsorption of biliary cholesterol.


### Ezetimibe/statin combination therapy

The combination of ezetimibe and statin therapy may synergistically decrease cholesterol absorption and synthesis, since the effective inhibition of cholesterol synthesis and the subsequent reduction in serum cholesterol levels by statins lead to increases in serum plant-sterol levels, probably as a result of reduced biliary secretion and the enhanced absorption of these sterols [9].

In 769 patients with hypercholesterolemia, ongoing statin therapy plus ezetimibe led to significant changes in LDL cholesterol (of  $-25.1\%$ ) (HDL cholesterol  $+2.7\%$ ; TG  $-14.0\%$ ) compared to the results with placebo (LDL cholesterol  $-3.7\%$ , HDL cholesterol  $+1.0\%$ , and TG  $-2.9\%$ ) [10]. Among patients who were not at their LDL

cholesterol target, 71.5% of those who received statin plus ezetimibe vs. 18.9% of those who received statin plus placebo reached their target levels at the end-point. In multicenter, randomized, double-blind, placebo-controlled studies, an ezetimibe + simvastatin group showed significantly improved LDL cholesterol, HDL cholesterol, and TG compared to a simvastatin-alone group. Ezetimibe plus simvastatin provided an incremental 13.8% LDL cholesterol decrease, 2.4% HDL cholesterol increase, and 7.5% TG decrease compared to simvastatin alone (Fig. 3) [6]. We also performed the Fluvastatin Add-On to Ezetimibe Trial (FAET) in patients with hypercholesterolemia (Fig. 4). After 6 months of ezetimibe monotherapy (10 mg/day), plasma LDL cholesterol decreased by 40 mg/dL (mean: 175–135 mg/dL at 6 months). Ezetimibe plus low-dose of fluvastatin (20 mg/day) combination therapy was started in poor-responders (LDL cholesterol:  $158 \pm 4$  mg/dL after 6 months ezetimibe monotherapy), then the LDL cholesterol levels reduced to  $130 \pm 4$  mg/dL ( $-28$  mg/dL), while LDL cholesterol in ezetimibe monotherapy group (responders) was  $128 \pm 4$  mg/dL. No adverse effect was observed. Although coadministration of ezetimibe plus low-dose of fluvastatin was more effective in reducing plasma concentrations of LDL-C than ezetimibe alone, we need to analyze the patient's characteristics of poor-responder to ezetimibe.

Combination therapy with 10 mg/day ezetimibe and 10 mg/day atorvastatin for 8 weeks



**Figure 4** Changes in serum LDL cholesterol levels in patients with hypercholesterolemia ( $n = 59$ ) (Fluvastatin Add-On to Ezetimibe Trial). After 6 months of ezetimibe monotherapy (solid line), combination therapy (ezetimibe + fluvastatin, dotted line) was started in poor-responders ( $n = 26$ ) after 6 months ezetimibe monotherapy. Responders ( $n = 33$ ) continued ezetimibe monotherapy (gray line).

resulted in significantly decreased total serum cholesterol and TG compared to 40 mg/day atorvastatin alone [11]. Ezetimibe/atorvastatin therapy significantly improved LDL cholesterol, HDL cholesterol, TG and high-sensitivity C-reactive protein (hs-CRP) compared to atorvastatin alone [12]. The addition of ezetimibe to on-going simvastatin treatment also resulted in a significantly greater percent reduction in LDL cholesterol from baseline (25.2%) compared to the results with placebo (0.9%). In a double-blind, multicenter, 6-week, parallel-group study, hypercholesterolemic patients were randomized to receive ezetimibe/simvastatin or rosuvastatin. At all doses and across doses, ezetimibe/simvastatin reduced LDL cholesterol significantly more than rosuvastatin [13]. Ezetimibe/simvastatin also produced significantly greater reductions in total cholesterol, non-HDL cholesterol and apolipoprotein B. Intensive lipid-lowering therapy with rosuvastatin 40 mg/day provided a greater LDL cholesterol-lowering effect than atorvastatin 80 mg/day, which enabled more patients to achieve their LDL cholesterol goals [13]. The EXPLORER study was designed to investigate the efficacy and safety of rosuvastatin 40 mg/day alone or in combination with ezetimibe 10 mg/day in patients with a high risk of coronary heart disease [14]. Interestingly, the combination of rosuvastatin/ezetimibe reduced LDL cholesterol significantly more than rosuvastatin alone. Other components of the lipid/lipoprotein profile were also significantly improved with rosuvastatin/ezetimibe. This combination therapy may have the strongest lipid-lowering effect.

The combination of ezetimibe/statin therapy reduced LDL cholesterol by 57% in familial hypercholesterolemia (FH) patients, regardless of the type of LDL receptor mutation. In clinical trials, ezetimibe/simvastatin produced greater reductions in LDL cholesterol than did monotherapy [15]. A possible molecular mechanism of the enhanced efficacy of ezetimibe plus simvastatin is decreased very (V) LDL and LDL apoB-100 concentrations through reduced VLDL production and the upregulation of LDL receptor-mediated LDL clearance. The additional decrease in plasma LDL cholesterol induced by ezetimibe showed wide inter-individual variability and was negatively correlated with the percent decrease in LDL cholesterol due to statin alone [16]. The reduction in LDL cholesterol in FH heterozygotes following combined therapy may affect the complex interplay between hepatic synthesis and the intestinal absorption of cholesterol. The usefulness of ezetimibe in enhancing the efficacy of apheresis therapy was evaluated in six Japanese patients with homozygous FH undergoing

LDL-apheresis in combination with atorvastatin or simvastatin [17]. Thus ezetimibe also appears to be useful in combination with statins for FH patients.

Since diabetes mellitus (DM) patients have been shown to have more NPC1L1 mRNA than control subjects [18], ezetimibe may be useful for the treatment of patients with DM in whom cholesterol absorption may be upregulated. In patients with type 2 DM, combination therapy is usually required to optimize glucose metabolism as well as to help patients achieve aggressive targets for LDL cholesterol and other lipid parameters associated with cardiovascular risk. Ezetimibe/simvastatin therapy was effective and well tolerated under actual practice conditions in high-risk patients with coronary heart disease and/or DM [19]. In a randomized, double-blind, multicenter study in type 2 DM patients, LDL cholesterol was reduced significantly more by adding ezetimibe 10 mg/day to simvastatin 20 mg/day than by doubling the dose of simvastatin to 40 mg/day [20]. In addition, ezetimibe + simvastatin therapy also produced significant incremental reductions in non-HDL cholesterol, very LDL cholesterol and apolipoprotein B relative to simvastatin 40 mg/day.

### **Ezetimibe or ezetimibe/statin in combination with fibrate therapy**

Although fenofibrate did not significantly reduce the risk of the primary outcome of coronary events, it did reduce total cardiovascular events, mainly due to fewer non-fatal myocardial infarctions and revascularizations in patients with type 2 DM [21]. Mixed hyperlipidemia is an important risk factor for the development of cardiovascular disease. Fenofibrate and ezetimibe offer complementary benefits to the lipid profile in patients with mixed hyperlipidemia. Patients in the fenofibrate plus ezetimibe and fenofibrate groups continued on their respective base study treatment, and patients in the ezetimibe and placebo groups were switched to fenofibrate plus ezetimibe and fenofibrate, respectively [22]. Fenofibrate plus ezetimibe produced significantly greater reductions in LDL cholesterol than fenofibrate alone. There were also significantly greater improvements in TG, HDL cholesterol, total cholesterol, non-HDL cholesterol, and apolipoprotein B with fenofibrate plus ezetimibe than with fenofibrate alone. In a multicenter, randomized, double-blind, placebo-controlled, parallel arm trial, the LDL cholesterol level was significantly reduced with ezetimibe/simvastatin + fenofibrate compared fenofibrate or placebo, but not compared

to ezetimibe/simvastatin [23]. HDL cholesterol and apolipoprotein A-I levels were significantly increased with ezetimibe/simvastatin + fenofibrate treatment compared to with ezetimibe/simvastatin or placebo. TG, non-HDL cholesterol, and apolipoprotein B levels were significantly reduced with ezetimibe/simvastatin + fenofibrate vs. all other treatments. Thus, the coadministration of ezetimibe/statin plus fenofibrate effectively improved the overall atherogenic lipid profile of patients with hyperlipidemia.

## Genetic studies on ezetimibe therapy

The NPC1L1 gene may have naturally occurring coding mutations. These mutations might affect the response to ezetimibe among individuals who carry such mutants. In fact, three common single-nucleotide polymorphisms have been found in NPC1L1 (1735C > G [L272L], 25342A > C and 27677T > C [V1296]). Different NPC1L1 protein variants, such as Val<sup>55</sup> to Leu<sup>55</sup> and Ile<sup>1233</sup> to Asp<sup>1233</sup> have been found in a non-responder to ezetimibe [24]. In addition, 101 patients with primary hypercholesterolemia had a significantly greater reduction in plasma LDL cholesterol with ezetimibe than subjects with at least one copy of the haplotype. A detailed characterization of DNA variations in NPC1L1, the target of ezetimibe, demonstrated that common variants in this gene are significantly associated with the LDL cholesterol reduction in response to treatment with ezetimibe combined with a statin, but not with baseline LDL cholesterol [25]. Moreover, genetic variation in NPC1L1 contributes to the variability in cholesterol absorption and plasma levels of LDL cholesterol [26]. Rare variants identified in low-absorbers were found in 6% of 1832 African-Americans and were associated with lower plasma levels of LDL cholesterol. Although none of these common or rare variants have been confirmed, evidence has indicated an association between LDL cholesterol and NPC1L1 gene production as a target for ezetimibe.

## Safety of ezetimibe

In 2382 patients with primary hypercholesterolemia, the coadministration of ezetimibe plus statin was well-tolerated [27]. In addition, in a randomized, double-blind, placebo-controlled trial, the safety and tolerability profiles for the ezetimibe/simvastatin and ezetimibe monotherapy groups were similar [28]. Ezetimibe plus statin therapy was well-tolerated and had a favorable

safety profile in both male and female subgroups [29]. The coadministration of ezetimibe and statin was equally safe across specific age groups: age <65 vs. ≥65 years; age <75 vs. more than 75 years [30]. Simvastatin at doses up to 40 mg is a well-tolerated and effective therapy for children with heterozygous familial hypercholesterolemia [31]. In addition, the combination of fenofibrate plus ezetimibe was well tolerated during the extension study, with similar rates of ALT/AST elevations between fenofibrate plus ezetimibe and fenofibrate alone [22]. Overall, ezetimibe shows a favorable safety profile across all patients.

## Conclusions

Ezetimibe is one of the most important medications for inhibiting cholesterol absorption. It has been shown to significantly reduce LDL cholesterol and improve other lipid parameters as either monotherapy or in combination therapy with other lipid-lowering medications, such as statins and fibrates. Such combination therapy is a more effective therapeutic option for the intensive lowering of LDL cholesterol. In addition, this therapy has been shown to have a favorable safety profile across all patients. Large randomized trials will be needed to evaluate the effectiveness of ezetimibe/statin or ezetimibe/fibrate therapy on clinical endpoints for the risk of coronary artery disease.

## References

- [1] Clader JW. The discovery of ezetimibe: a view from outside the receptor. *J Med Chem* 2004;47:1–9.
- [2] Strandberg TE, Tilvis RS, Pitkala KH, Miettinen TA. Cholesterol and glucose metabolism and recurrent cardiovascular events among the elderly: a prospective study. *J Am Coll Cardiol* 2006;48:708–14.
- [3] Altmann SW, Davis Jr HR, Zhu LJ, Yao X, Hoos LM, Tetzloff G, et al. Niemann-Pick C1 Like 1 protein is critical for intestinal cholesterol absorption. *Science* 2004;303:1201–4.
- [4] Garcia-Calvo M, Lisnock J, Bull HG, Hawes BE, Burnett DA, Braun MP, et al. The target of ezetimibe is Niemann-Pick C1-Like 1 (NPC1L1). *Proc Natl Acad Sci USA* 2005;102:8132–7.
- [5] van Heek M, Farley C, Compton DS, Hoos L, Alton KB, Sybertz EJ, et al. Comparison of the activity and disposition of the novel cholesterol absorption inhibitor, SCH58235, and its glucuronide, SCH60663. *Br J Pharmacol* 2000;129:1748–54.
- [6] Davidson MH, McGarry T, Bettis R, Melani L, Lipka LJ, LeBeaut AP, et al. Ezetimibe coadministered with simvastatin in patients with primary hypercholesterolemia. *J Am Coll Cardiol* 2002;40:2125–34.
- [7] Kalogirou M, Tsimihodimos V, Gazi I, Filippatos T, Saougos V, Tselepis AD, et al. Effect of ezetimibe monotherapy on the concentration of lipoprotein subfractions in



- patients with primary dyslipidaemia. *Curr Med Res Opin* 2007;23:1169–76.
- [8] Geiss HC, Otto C, Parhofer KG. Effect of ezetimibe on low-density lipoprotein subtype distribution: results of a placebo-controlled, double-blind trial in patients treated by regular low-density lipoprotein apheresis and statins. *Metabolism* 2006;55:599–604.
  - [9] Miettinen TA, Gylling H, Lindholm N, Miettinen TE, Rajaratnam RA, Relas H, et al. Serum noncholesterol sterols during inhibition of cholesterol synthesis by statins. *J Lab Clin Med* 2003;141:131–7.
  - [10] Gagné C, Bays HE, Weiss SR, Mata P, Quinto K, Melino M, et al. Efficacy and safety of ezetimibe added to ongoing statin therapy for treatment of patients with primary hypercholesterolemia. *Am J Cardiol* 2002;90:1084–91.
  - [11] Bulut D, Hanefeld C, Bulut-Streich N, Graf C, Mügge A, Spiecker M. Endothelial function in the forearm circulation of patients with the metabolic syndrome—effect of different lipid-lowering regimens. *Cardiology* 2005;104:176–80.
  - [12] Ballantyne CM, Houry J, Notarbartolo A, Melani L, Lipka LJ, Suresh R, et al. Effect of ezetimibe coadministered with atorvastatin in 628 patients with primary hypercholesterolemia: a prospective, randomized, double-blind trial. *Circulation* 2003;107:2409–15.
  - [13] Catapano AL, Davidson MH, Ballantyne CM, Brady WE, Gazzara RA, Tomassini JE, et al. Lipid-altering efficacy of the ezetimibe/simvastatin single tablet versus rosuvastatin in hypercholesterolemic patients. *Curr Med Res Opin* 2006;22:2041–53.
  - [14] Ballantyne CM, Weiss R, Moccetti T, Vogt A, Eber B, Sosef F, et al. Efficacy and safety of rosuvastatin 40mg alone or in combination with ezetimibe in patients at high risk of cardiovascular disease (results from the EXPLORER study). *Am J Cardiol* 2007;99:673–80.
  - [15] Telford DE, Sutherland BG, Edwards JY, Andrews JD, Barrett PH, Huff MW. The molecular mechanisms underlying the reduction of LDL apoB-100 by ezetimibe plus simvastatin. *J Lipid Res* 2007;48:699–708.
  - [16] Pisciotto L, Fasano T, Bellocchio A, Bocchi L, Sallo R, Fresa R, et al. Effect of ezetimibe coadministered with statins in genotype-confirmed heterozygous FH patients. *Atherosclerosis* 2007;194:e116–22, 2005;366:1849–61.
  - [17] Yamamoto A, Harada-Shiba M, Endo M, Kusakabe N, Tanioaka T, Kato H, et al. The effect of ezetimibe on serum lipids and lipoproteins in patients with homozygous familial hypercholesterolemia undergoing LDL-apheresis therapy. *Atherosclerosis* 2006;186:126–31.
  - [18] Lally S, Tan CY, Owens D, Tomkin GH. Messenger RNA levels of genes involved in dysregulation of postprandial lipoproteins in type 2 diabetes: the role of Niemann-Pick C1-like 1, ATP-binding cassette, transporters G5 and G8, and of microsomal triglyceride transfer protein. *Diabetologia* 2006;49:1008–16.
  - [19] Hildemann SK, Barho C, Karmann B, Darius H, Bestehorn K. Dual cholesterol inhibition with ezetimibe/simvastatin in pre-treated hypercholesterolaemic patients with coronary heart disease or diabetes mellitus: prospective observational cohort studies in clinical practice. *Curr Med Res Opin* 2007;23:713–9.
  - [20] Gaudiani LM, Lewin A, Meneghini L, Perevozskaya I, Plotkin D, Mitchel Y, et al. Efficacy and safety of ezetimibe co-administered with simvastatin in thiazolidinedione-treated type 2 diabetic patients. *Diabetes Obes Metab* 2005;7:88–97.
  - [21] Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, et al. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2007;370:1687–97.
  - [22] McKenney JM, Farnier M, Lo KW, Bays HE, Perevozskaya I, Carlson G, et al. Safety and efficacy of long-term co-administration of fenofibrate and ezetimibe in patients with mixed hyperlipidemia. *J Am Coll Cardiol* 2006;47:1584–7.
  - [23] Farnier M, Roth E, Gil-Extremiera B, Mendez GF, Macdonell G, Hamlin C, et al. Efficacy and safety of the coadministration of ezetimibe/simvastatin with fenofibrate in patients with mixed hyperlipidemia. *Am Heart J* 2007;153:335.e1–8.
  - [24] Wang J, Williams CM, Hegele RA. Compound heterozygosity for two non-synonymous polymorphisms in NPC1L1 in a non-responder to ezetimibe. *Clin Genet* 2005;67:175–7.
  - [25] Simon JS, Karnoub MC, Devlin DJ, Arreaza MG, Qiu P, Monks SA, et al. Sequence variation in NPC1L1 and association with improved LDL-cholesterol lowering in response to ezetimibe treatment. *Genomics* 2005;86:648–56.
  - [26] Cohen JC, Pertsemlidis A, Fahmi S, Esmail S, Vega GL, Grundy SM, et al. Multiple rare variants in NPC1L1 associated with reduced sterol absorption and plasma low-density lipoprotein levels. *Proc Natl Acad Sci USA* 2006;103:1810–5.
  - [27] Davidson MH, Ballantyne CM, Kerzner B, Melani L, Sager PT, Lipka L, et al. Efficacy and safety of ezetimibe coadministered with statins: randomised, placebo-controlled, blinded experience in 2382 patients with primary hypercholesterolemia. *Int J Clin Pract* 2004;58:746–55.
  - [28] Goldberg AC, Sapre A, Liu J, Capece R, Mitchel YB, Ezetimibe Study Group. Efficacy and safety of ezetimibe coadministered with simvastatin in patients with primary hypercholesterolemia: a randomized, double-blind, placebo-controlled trial. *Mayo Clin Proc* 2004;79:620–9.
  - [29] Bennett S, Sager P, Lipka L, Melani L, Suresh R, Veltri E, et al. Consistency in efficacy and safety of ezetimibe coadministered with statins for treatment of hypercholesterolemia in women and men. *J Womens Health* 2004;13:1101–7.
  - [30] Lipka L, Sager P, Strony J, Yang B, Suresh R, Veltri E, et al. Efficacy and safety of coadministration of ezetimibe and statins in elderly patients with primary hypercholesterolaemia. *Drugs Aging* 2004;21:1025–32.
  - [31] de Jongh S, Ose L, Szamosi T, Gagné C, Lambert M, Scott R, et al. Efficacy and safety of statin therapy in children with familial hypercholesterolemia: a randomized, double-blind, placebo-controlled trial with simvastatin. *Circulation* 2002;106:2231–7.

